The composition of matter of claim 67 in which said plurality of sub-populations is comprised of one of said sub-populations for each charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

76. The composition of matter of claim 67 in which said populations of multiply charged ions is formed by:

14

5

dispersing a solution containing said polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field; and

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said distinct polyatomic parent molecular species becomes dispersed in said bath gas as said multiply charged ions.

The composition of matter of claim of in which said single polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

72. The composition of matter of claim of in which said single polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

73. The composition of matter of claim 57 in which a mass spectrum is generated from said population of multiply charged ions, said mass spectrum comprising a sequence of peaks due to said plurality of sub-populations of said population of multiply charged ions and analyzing said peaks of said mass spectrum to determine a value of said molecular weight of said single polyatomic parent molecular species.

The composition of matter as claimed in claim of in which said single polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

A composition of matter comprising a population of multiply charged ions derived from a sample substantially comprising a single polyatomic parent molecular species, the number of charges on each ion defining said ion's charge state number, said population of said multiply charged ions comprising a plurality of sub-populations, the ions of a distinct sub-population having the same charge number which differs from said charge number of an adjacent said sub-population by one, each of said sub-populations having said charge state number not less than five, said composition of matter formed by:

dispersing a solution containing said polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field; and

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said polyatomic parent molecular species become dispersed in said bath gas as said multiply charged ions.

76. The composition of matter of claim 75 in which said charge state number of each of said sub-populations is not less than seven.

10

-17

7

The composition of matter of claim 25 in which all molecules of said single polyatomic parent molecular species have substantially the same molecular weight.

78. The composition of matter of claim 75 in which said single polyatomic parent molecular species is selected from the class of compounds known as biopolymers.

9

The composition of matter of claim \mathcal{B} such that when a mass spectrum is generated from said population of multiply charged ions, said mass spectrum comprises a sequence of peaks due to said plurality of sub-populations of said population of multiply charged ions and analyzation of said peaks of said mass spectrum determines a value of the molecular weight of said single polyatomic parent molecular species.

15

20. The composition of matter of claim 75 in which said single polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

1:14

281. The composition of matter of claim 79 in which said single polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

1:16

charged ions generated from a sample comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a plurality of sub-populations, and for each of said populations, there being one of said sub-populations for each possible charge state number whose minimum value is not less than three and whose maximum value is not less than five

:17

83. The composition of matter of claim 82 in which said plurality of sub-populations for each of said populations is comprised of one of said sub-populations for each charge state number whose minimum value is not less than five and whose maximum value is not less than seven.

18

At. The composition of matter of claim 22 in which said plurality of sub-populations for each of said populations is comprised of one of said sub-populations for each charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

The composition of matter of claim 22 in which all molecules of each of said distinct polyatomic parent molecular species have substantially the same molecular weight.

16

26. The composition of matter of claim 22 in which at least one of said distinct polyatomià parent molecular species is selected from a class of compounds known as biopolymers.

distinct 31. The composition of matter of claim 32 in which at least one of said, discrete polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

28. The composition of matter of claim 22 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

20

21

D

D

D

89. The composition of matter of claim 82 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

98. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a plurality of subpopulations, the ions of each sub-population within said plurality having the same charge state number which differs from said charge state number of another of said sub-populations within said plurality, each of said sub-populations having said charge state number not less than five.

The composition of matter of claim of in which each of said sub-populations has said charge state number not less than seven.

92. The composition of matter of claim 90 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

93. The composition of matter of claim 90 in which at least one of said discrete polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

D

D

D

94. The composition of matter of claim 90 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

95. The composition of matter of claim on which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

30
96. The composition of matter of claim 90 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 7000 amu.

A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a plurality of sub-populations, and for each of said populations, there being one of said sub-populations are each possible charge state number whose minimum value is not less than three and

whose maximum value is not less than five, said composition of matter being formed by:

dispersing a solution containing said one or more distinct polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field; and

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said distinct polyatomic parent molecular species become dispersed in said bath gas as said multiply charged ions.

98. The composition of matter of claim 97 in which said plurality of sub-populations; for each of said populations, is comprised of one of said sub-populations for each charge state number whose minimum value is not less than five and whose maximum value is not less than seven.

99. The composition of matter of claim 97 in which said plurality of sub-populations, for each of said populations is comprised of one of said sub-populations for each charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

D

D

D

100. The composition of matter of claim of in which all molecules of a given of said distinct polyatomic parent molecular species have substantially the same molecular weight.

35
101. The composition of matter of claim 97 in which all molecules of at least one of said distinct polyatomic parent molecular species have the same chemical formula.

36. The composition of matter of claim 37 in which all molecules of each of said distinct polyatomic parent molecular species are chemically indistinct.

37
103. The composition of matter of claim 97 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

36
31
104. The composition of matter of claim 97 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

105. The composition of matter of claim 97 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

40 31 106. The composition of matter of claim of in which said bath gas is heated.

107. The composition of matter of claim 97 in which said dispersing of said charged droplets is substantially in a first direction and said bath gas is directed in a substantially counter current direction to said first direction.

108. The composition of matter of claim 97 in which said bath gas is at approximately atmospheric pressure.

109. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a plurality of subpopulations, and for each of said populations there being a one-of-said sub-populations

D

for each of every possible charge state number whose minimum value is not less than three and whose maximum value is not less than five, said composition of matter being useful for:

D

D

D

D

D

D

41

from the results of said mass analysis obtaining mass/charge (m/z)

from the results of said mass analysis obtaining mass/charge (m/z)

values for said ions of said sub-populations; and

determining a value of the molecular weight of said distinct polyatomic parent molecular species from the mass/charge (m/z) values of said ions in said sub-populations.

43

populations for each of said populations is comprised of one of said sub-populations for each charge state number whose minimum value is not less than five and whose maximum value is not less than seven.

J.H. The composition of matter of claim 100 in which said plurality of subpopulations for each of said populations is comprised of one of said sub-populations
for each charge state number whose minimum value is not less than seven and whose
maximum value is not less than ten.

142. The composition of matter of claim 199 in which all molecules of a given of said distinct polyatomic parent molecular species have substantially the same molecular weight.

113. The composition of matter of claim 109 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

The composition of matter of claim 109 in which at least one of said discrete polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

D

D-

D

D

143. The composition of matter of claim 109 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

116. The composition of matter of claim 109 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

15. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations comprising a plurality of sub-populations, the ions of a distinct sub-population within said plurality having the same charge number which differs from said charge number of another of said sub-populations within said plurality, each of said sub-populations having said charge state number not less than five; said composition of matter being useful for:

from the results of said mass analysis obtaining mass/charge (m/z)

from the results of said mass analysis obtaining mass/charge (m/z)

from the results of said mass analysis obtaining mass/charge (m/z)

values for said ions of said sub-populations; and

determining a value of the molecular weight of said distinct polyatomic parent molecular species from the mass/charge (m/z) values of said ions in said sub-populations.

The composition of matter of claim 117 in which each of said sub-populations has said charge state number not less than seven.

53
119. The composition of matter of claim 117 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

126. The composition of matter of claim 147 in which at least one of said discrete polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

D

D

D

D

: 57

The composition of matter of claim 117 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising polyethylene glycols.

122. The composition of matter of claim 117 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

123. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the number of charges on each ion defining the ion's charge state number, each of said populations comprising a plurality of sub-populations, and for each of said populations, there being one of said sub-populations for each possible charge state number whose minimum value is not less than three and whose maximum value is not less than five, said composition of matter being formed and mass-analyzed by:

bath gas as charged droplets, said dispersing taking place in the presence of an electric field;

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said distinct polyatomic parent molecular species becomes dispersed in said bath gas as said multiply charged ions;

D

ID

D

D D passing a portion of said multiply charged ions in said bath gas into vacuum containing a mass analyzer, carrying out a mass analysis of the ions in said one or more populations and from the results of said mass analysis obtaining mass/charge (m/z) values for said ions of said subfrom the results of said mass analysis populations; and

determining a value of the molecular weight of said distinct polyatomic parent molecular species from the mass/charge (m/z) values of said ions in said sub-populations.

124. The composition of matter of claim 123 in which said plurality of subpopulations, for each of said populations, is comprised of one of said sub-populations
for each charge state number whose minimum value is not less than five and whose
maximum value is not less than seven.

populations, for each of said populations, is comprised of one of said sub-populations for each charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

D

The composition of matter of claim 123 in which all molecules of a given of said distinct polyatomic parent molecular species have substantially the same molecular weight.

61

51 127. The composition of matter of claim 123 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as

62

biopolymers.

128. The composition of matter of claim 123 in which at least one of said distinct polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

63

129. The composition of matter of claim 123 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising synthetic polymers comprising less than four distinct elements, said group comprising polyethylene glycols.

57

64

130. The composition of matter of claim 123 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

134. The composition of matter of claim 123 in which said bath gas is heated.

ماما

132. The composition of matter of claim 123 in which said dispersing of said charged droplets is substantially in a first direction and said bath\gas is directed in a substantially counter current direction to said first direction,

67

133. The composition of matter of claim 123 in which said bath gas is at approximately atmospheric pressure.

134. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, said polyatomic parent molecular species not comprising polyethylene glycol, the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a plurality of sub-populations, and for each of said populations, there being one of said sub-populations for each possible charge state number whose minimum value is not less than three and whose maximum value is not less than five, the formation and mass analysis of said composition of matter being performed by:

dispersing a solution containing said polyatomic parent species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field;

allowing the solvent of said solution to evaporate from said charged droplets
until at least some molecules of said polyatomic parent molecular
species become dispersed in said bath gas as said multiply charged ions;

passing a portion of said multiply charged ions in said bath gas into vacuum containing a mass analyzer, carrying out a mass analysis of the ions in said one or more populations and from the results of said mass analysis obtaining mass/charge (m/z) values for said ions of said subfrom the results of Said mass analysis
populations; and

determining a value of the molecular weight of said distinct polyatomic parent molecular species from the mass/charge (m/z) values of said ions in said sub-populations.

135. The composition of matter of claim 134 in which said plurality of subpopulations, for each of said populations is comprised of one of said sub-populations

15

D D

Ch.

D

Þ

D

D

D

for every charge state number whose minimum value is not less than five and whose maximum value is not less than seven.

70 \ 68

D

D

D

D

D

٢),

D

D

D

populations, for each of said populations, is comprised of one of said sub-populations for each charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

137. The composition of matter of claim 134 in which all molecules of a given of said distinct polyatomic parent molecular species have substantially the same molecular weight.

138. The composition of matter of claim 134 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

139. The composition of matter of claim 134 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

140. A composition of matter derived from a sample comprising one or more distinct polyatomic parent molecular species, all molecules of each of said distinct polyatomic parent molecular species having substantially the same molecular weight, said of multiply charged from composition of matter comprising a population for each of said distinct polyatomic parent molecular species of multiply charged ions, the generation of a mass spectrum each of said distinct polyatomic parent molecular species of multiply charged ions, the generation of a mass spectrum each of said distinct polyatomic parent molecular species of not less than two peaks for each distinct population, each peak corresponding to a discrete subpopulation of multiply charged ions, each of said multiply charged ions having a discrete integral number of charges not less than five wherein said number of charges of the multiply charged ions of each peak in said sequence for each distinct

polyatomic parent molecular species differs from those of an adjacent peak by one charge unit.

75 141. The composition of matter of claim 140 in which at least one of said subpopulations\has said number of charges not less than seven.

142. The composition of matter of claim 140 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

.. 17

143. The composition of matter of claim 140 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising polyethylene glycols.

74 144. The composition of matter of claim 140 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 5000 amu.

79

145. The composition of matter of claim 140 in which at least one of said distinct polyatomic parent molecular species has a molecular weight not less than about 7000 amu.

74

ં વૂ ૦

146. A composition of matter comprising one or more distinct populations of multiply charged ions, each of said multiply charged ions in said distinct populations of multiply charged ions being eharacterizable by the symbol xi, the numerical value of xi being the m/z value for said one of said multiply charged ions such that xi = M/i + m_a wherein M_r is the molecular weight of a distinct parent polyatomic molecular species from which all of said multiply charged ions in said distinct population of multiply charged ions are derived, i is an integer equal to the number of adduct charges attached to (or removed from as in the case with negative ions) said distinct

D

D

D

parent polyatomic molecular species to form said multiply charged ions, m_a is the average mass of said individual adduct charges attached to (or-removed from) said multiply charged ions, each of said distinct population of ions comprising a plurality of sub-populations, the ions of each sub-population having the same values for i, m_a and M_r and therefore the same value of xi, said plurality of sub-populations comprising one sub-population for each integral value of i whose minimum is not less than three and whose maximum is not less than five.

D

D

D

D

D

96

147. The composition of matter of claim 146 in which there is at least one plurality of sub-populations comprised of one sub-population for each integral value of said i whose minimum is not less than five and whose maximum is not less than seven.

148. The composition of matter of claim 146 in which there is at least one plurality of sub-populations comprised of one sub-population for each integral value of said i whose minimum is not less than seven and whose maximum is not less than ten.

75. The composition of matter of claim 146 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

84 go destrict.

150. The composition of matter of claim 146 in which at least one of said discrete polyatomic parent molecular species is selected from the group comprising proteins, peptides, polypeptides, carbohydrates, oligonucleotides and glycoproteins.

151. The composition of matter of claim 146 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising polyethylene glycols.

152. The composition of matter of claim 146 characterized by, when a mass spectrum is generated from at least one of said distinct populations of multiply charged ions,

said mass spectrum comprises a sequence of peaks of at least one of said plurality of sub-populations of said distinct population of multiply charged ions and, upon analyzing said peaks of said mass spectrum, a determination can be made of the value of said molecular weight, M_r, of said distinct polyatomic parent molecular species.

D

D

D

D

D

D

153. The composition of matter of claim 146 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, M_r , not less than about 5000 arou.

154. The composition of matter of claim 146 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, M_r, not less than about 7000 amu.

charged ions, each of said multiply charged ions in said distinct populations of multiply charged ions being characterizable by the symbol xi, the numerical value of xi being the m/z value for one of said multiply charged ions such that xi = M_t/i + m_a wherein M_t is the molecular weight of a distinct parent polyatomic molecular species from which all of said multiply charged ions in said distinct population of multiply charged ions are derived, i is an integer equal to the number of adduct charges attached to (or removed from as in the case with negative ions) said distinct parent polyatomic molecular species to form said multiply charged ions, m_a is the average mass of said individual adduct charges attached to (or removed from) said multiply charged ion, each of said distinct population of ions comprising a plurality of subpopulations, the ions of each sub-population having the same values for i, m_a and M_t and therefore the same value of xi, said plurality of sub-populations comprising distinct sub-populations with distinct values of i, each of said sub-populations having a value for i not less than five.

D

D

91

95

D

The composition of matter of claim 155 in which each of at least one of said plurality of sub-populations is comprised of distinct sub-populations with distinct values of i, each of said sub-populations having a value for i not less than seven.

157. The composition of matter of claim 155 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

92

158. The composition of matter of claim 155 characterized in that when a mass spectrum is generated from at least one of said distinct populations of multiply charged ions, said mass spectrum comprises a sequence of peaks having at least one of said plurality of sub-populations of said distinct population of multiply charged ions and, when said mass spectrum is analyzed, said peaks determine a value of said molecular weight, M_r, of said distinct polyatomic parent molecular species.

93

159. The composition of matter of claim 155 in which at least one of said distinct polyatomic parent molecular species has a value of said molecular weight, M_r , not less than about 5000 amu.

160. The composition of matter of claim 155 in which at least one of said distinct polyatomic parent molecular species has a value of said molecular weight. Monot least one of said molecular weight.

polyatomic parent molecular species has a value of said molecular weight, M_r, not less than about 7000 amu.

161. A composition of matter comprising one or more distinct populations of multiply charged ions, each of said multiply charged ions in said distinct populations of multiply charged ions being characterized by the symbol xi, the numerical value of xi being the m/z value for said one of said multiply charged ions such that $xi = M_r/i + m_a$ wherein M_r is the molecular weight of a distinct parent polyatomic molecular species from which all of said multiply charged ions in said distinct population of multiply charged ions are derived, i is an integer equal to the number of adduct

charges attached to (or removed from as in the case with negative ions) said distinct parent molecular species to form said multiply charged ions, ma is the average mass of said individual adduct charges attached to (or removed from) said multiply charged populations ion, each of said distinct population of ions comprising a plurality of sub-populations, the ions of each sub-population having the same values for i, ma and Mr and therefore the same value of xi, said plurality of sub-populations comprising at least one subpopulation for each integral value of i whose minimum is not less than three and whose maximum ix not less than five, said composition of matter being formed by:

D

D

D

D

D

D

D

D

97

98

dispersing a solution containing said polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of a electric field; and

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said parent polyatomic molecular species become dispersed in said bath gas as said multiply charged ions.

162. The composition of matter of claim 161 in which there is at least one of said plurality of sub-populations, comprised of one sub-population, for each integral value of said i whose minimum is not less than five and whose maximum is not less than seven.

95 163. The composition of matter of claim 164 in which there is at least one of said plurality of sub-populations comprised of one sub-population, for each integral value of said i whose minimum is not less than seven and whose maximum is not less than ten.

164. The composition of matter of claim 161 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

The composition of matter of claim 161 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising polyethylene glycols.

100

166. The composition of matter of claim 161 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, M_r, not less than about 5000 amu.

101

D

D

D

D

D

D

D

167. A composition of matter comprising one or more distinct populations of multiply charged ions, each of said multiply charged ions in said distinct populations of multiply charged ions being characterizable by the symbol xi, the numerical value of xi being the m/z value for said one of said multiply charged ions such that xi = M/i + m_a wherein M_r is the molecular weight of a distinct parent polyatomic molecular species from which all of said multiply charged ions in said distinct population of multiply charged ions are derived, i is integer equal to the number of adduct charges attached to (or removed from as in the case with negative ions) said distinct parent polyatomic molecular species/to form said multiply charged ions, m_a is the average mass of said individual adduct charges attached to (or removed from) said multiply charged ion, each of said distinct population of ions comprising a plurality of sub-populations, the ions of each sub-population having the same values for i, ma and M_r and therefore the same value of xi, said plurality of sub-populations comprising at least one sub-population for each integral value of in whose minimum is not less than three and whose maximum is not less than five, said composition of matter being characterized by the generation of a mass spectrum from at least one of said distinct populations of multiply charged ions comprising a sequence of peaks having at least one of said plurality of sub-populations of said distinct population of multiply charged ions and upon analyzing said peaks of said mass spectrum, a determination can be -made-of a value of said molecular weight, M_r, of said distinct polyatomic parent molecular species.

167 The composition of matter of claim 166

D

D

The composition of matter of claim 167 in which there is at least one of said plurality of sub-populations comprised of one sub-population for each integral value of said i whose minimum is not less than five and whose maximum is not less than seven.

103

169. The composition of matter of claim 167 in which there is at least one of said plurality of sub-populations comprised of one sub-population for each integral value of said i whose minimum is not less than seven and whose maximum is not less than ten.

170. The composition of matter of claim 167 in which said populations of multiply charged ions are formed by:

dispersing a solution containing said polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field; and

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said one or more distinct polyatomic parent molecular species become dispersed in said bath gas as said multiply charged ions.

105
177. The composition of matter of claim 167 in which at least one of said distinct polyatomic parent molecular species is selected from a class of compounds known as biopolymers.

172. The composition of matter of claim 167 in which at least one of said distinct polyatomic parent molecular species is not selected from the group comprising polyethylene glycols.

101

107 The composition of matter of claim 167 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, Mr, not less than about 5000 amu.

108 101

D

0

D

D

D

D

D

D

174. The composition of matter of claim 167 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, M_r, no less than about 7000 arru.

109 175. A composition of matter comprising one or more distinct populations of multiply charged ions, each of said\multiply charged ions in said distinct populations of multiply charged ions being characterizable by the symbol xi, the numerical value of xi being the m/z value for-said one of said multiply charged ions such that $xi = M_r/i + m_a$ wherein M_r is the molecular weight of a distinct parent polyatomic molecular species from which all of said multiply charged ions in said distinct population of multiply charged ions are derived, i is an integer equal to the number of adduct charges attached to (or removed from as in the case with negative ions) said distinct polyatomic parent molecular species to form said multiply charged ions, ψ_a is the average mass of said individual adduct charges attached to (or removed from) said multiply charged ion, each of said distinct population of ions comprising a plurality of sub-populations, the ions of each subpopulations having the same values for i, m, and M, and therefore the same value of xi, said plurality of sub-populations comprising distinct sub-populations with distinct values of i, each of said sub-populations having a value for i not less than five, said composition of matter being characterized such that generation of a mass spectrum from at least one of said distinct populations of multiply charged ions comprises à sequence of peaks due to at least one of said plurality of sub-populations of said distinct population of multiply charged ions and upon analyzing the same, said peaks of said mass spectrum allowed for the determination of a value of said molecular weight, M_r, of said distinct polyatomic parent molecular species, said composition of matter formed and mass analyzed by:

dispersing a solution containing said polyatomic parent species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field;

allowing the solvent of said solution to evaporate from said charged droplets until at least\some molecules of said polyatomic parent molecular species become dispersed in said bath gas as said multiply charged ions;

passing a portion of said multiply charged ions in said bath gas into vacuum containing a mass analyzer;

D

D

D

D

D

D

. 111

generation of said mass spectrum comprising a sequence of peaks due to said subpopulations of said one or more distinct populations; and

determination of the molecular weight of said one or more of said distinct polyatomic parent molecular species.

110 The composition of matter of claim which each of at least one plurality of sub-populations is comprised of distinct sub-populations with distinct values of, i, each of said sub-populations having a value for i not less than seven.

109 177. The composition of matter of claim 175 in which at least one of said distinct polyatomic parent molecular species has a value for said molecular weight, Mr, not less than about 5000 amu.

:112 109 178. The composition of matter of claim 175 in which at least one of said distinct polyatomic parent molecular species has a value, for said molecular weight, M_r, not less than about 7000 amu.

A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the molecular weight of said distinct polyatomic parent molecular species being at least about 5000 atomic mass units (amu), the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising a set of one or more sub-populations, the ions of a distinct sub-population within said set of sub-populations having the same charge state number which differs from said charge state number of another of said sub-populations within said set, each said sub-population having said charge state number not less than five, said composition of matter being characterized by:

D

114

116

D

D

D

D

P

carrying out a mass analysis of the ions in said one or more populations and from

the results of said mass analysis obtaining mass/charge (m/z) values for said

from the results of fact mass analysis

ions of said sub-populations; and

determining a value of the molecular weight of said distinct polyatomic parent molecular species from the mass/charge (m/z) values of said ions in said sub-populations.

180. The composition of matter of claim 179 in which each of at least one of said plurality of sub-populations is comprised of distinct sub-populations with distinct values of i, each of said sub-population having a value for i not less than seven.

distinct

181. The composition of matter of claim 179 in which all molecules of said single polyatomic parent molecular species have substantially the same molecular weight.

182. The composition of matter of claim 179 in which said single polyatomic parent molecular species is selected from the class of compounds known as biopolymers.

113 183. The composition of matter of claim 179 in which all molecules of at least one of said distinct polyatomic parent molecular species have the same chemical formula.

118

Þ

D

D

V

184. A composition of matter comprising one or more distinct populations of multiply charged ions generated from a sample comprising one or more distinct polyatomic parent molecular species, the molecular weight of said distinct polyatomic parent molecular species being at least about 5000 atomic mass units (amu), the number of charges on each ion defining the ion's charge state number, each of said populations of ions comprising one or more sub-populations, and for each of said populations, there being one of said sub-populations for each possible charge state number whose minimum value is not less than three and whose maximum value is not less than five, formation and mass analysis of said composition of matter being characterized by:

dispersing a solution containing said one or more distinct polyatomic parent molecular species into a bath gas as charged droplets, said dispersing taking place in the presence of an electric field;

allowing the solvent of said solution to evaporate from said charged droplets until at least some molecules of said distinct polyatomic parent molecular species become dispersed in said bath gas as said multiply charged ions;

passing a portion of said multiply charged ions in said bath gas into vacuum containing a mass analyzer;

generation of a mass spectrum comprising a sequence of peaks due to said subpopulations of said one or more distinct populations; and

the determination of the molecular weight of said one or more of said distinct polyatomic parent molecular species resulting in a value for the molecular weight.

185. The composition of matter of claim 184 in which said plurality of sub-populations, for each of said populations, is comprised of one of said sub-populations for each of every charge state number whose minimum value is not less than five and whose maximum value is not less than seven.

186. The composition of matter of claim 184 in which said plurality of sub-populations, for each of said populations, is comprised of one of said sub-populations for each of charge state number whose minimum value is not less than seven and whose maximum value is not less than ten.

187. The composition of the matter of claim 183 in which all molecules of a given of said distinct polyatomic parent molecular species have substantially the same molecular weight.